

Clenpiq Instructions

- To ensure a successful exam be sure to follow instructions Carefully.
- The entire day before your Colonoscopy be sure to follow a clear liquid diet.
- Do not drink any milk, anything colored red or purple, and no alcoholic beverages.
- You must have a ride for the day of your exam. If you fail to have acceptable transportation your procedure will be cancelled. Uber or Taxi are not acceptable modes of transportation, unless you have someone accompanying you.
- Let Dr. Adeyefa and his staff know if you are taking any blood thinners, such as Aspirin, Warfarin, Coumadin, Pradaxa, Effient, or Lovonox, as you may need to hold these medications before the procedure.
- If you are diabetic, take half your insulin dose the day of the procedure.
- If you take high blood pressure medication, take the medications the day of the procedure.
- If you take NSAID's such as Advil, Motrin, Celebrex, or ibuprofen you should discuss this with Dr. Adeyefa in advance.

At 5:00 pm the evening before the procedure:



Drink 1 bottle of Clenpiq prep with 5 8oz cups of clear liquid to stay hydrated. Do not rush through drinking all of this, drinking too fast will cause nausea and possible vomiting.

At 12:00 am (Midnight) the morning of the procedure:



Drink 1 bottle of Clenpiq prep with 5 8oz cups of clear liquid to stay hydrated. Do not rush through drinking all of this, drinking too fast will cause nausea and possible vomiting.